VOL. 535, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY DECEMBER 28, 1990

CONTENTS

14TH INTERNATIONAL SYMPOSIUM ON COLUMN LIQUID CHROMATOGRAPHY, BOSTON, MA, MAY 20–25, 1990, PART I

Foreword
by B. L. Karger 1

Key moments in the evolution of liquid chromatography
by L. S. Ettre (Norwalk, CT, U.S.A.) 3

Performance of wide-pore silica- and polymer-based packing materials in polypeptide separation: effect of pore size and alkyl chain length
by N. Tanaka, K. Kimata, Y. Mikawa, K. Hosoya and T. Araki (Kyoto, Japan), Y. Ohtsu and Y. Shiojima (Yokohama, Japan), R. Tsuboi (Kyoto, Japan) and H. Tsuchiya (Ibaraki, Japan) 13

Post-column continuous-flow analysis combined with reversed-phase liquid chromatography and computer-aided detection for the characterisation of peptides
by A. F. Fell and J. B. Castledine (Bradford, U.K.), B. Sellberg and R. Modin (Uppsala, Sweden) and R. Weinberger (Ramsey, NJ, U.S.A.) 33

Post-column reaction detection and flow injection analysis
by H. Engelhardt, R. Klinkner and G. Schöndorf (Saarbrücken, F.R.G.) 41

High-performance liquid chromatographic computer simulation based on a restricted multi-parameter approach. I. Theory and verification
by J. W. Dolan, D. C. Lommen and L. R. Snyder (Lafayette, CA, U.S.A.) 55

High-performance liquid chromatographic computer simulation based on a restricted multi-parameter approach. II. Applications
by L. R. Snyder, J. W. Dolan and D. C. Lommen (Lafayette, CA, U.S.A.) 75

Studies on system peaks in ion-pair adsorption chromatography. IV. Optimization of peak compression
by T. Fornstedt, D. Westerlund and A. Sokolowski (Uppsala, Sweden) 93

Porous zirconia and titania as packing materials for high-performance liquid chromatography
by U. Trüdinger, G. Müller and K. K. Unger (Mainz, F.R.G.) 111

New family of high-resolution ion exchangers for protein and nucleic acid purifications from laboratory to process scales
by D. M. Dion, K. O'Connor, D. Phillips, G. J. Vella and W. Warren (Milford, MA, U.S.A.) 127

Monosized stationary phases for chromatography
by T. Ellingsen, O. Aune and J. Ugelstad (Trondheim, Norway) and S. Hagen (Lillestrøm, Norway) 147

Synthesis of mixed-functional-phase silica supports for liquid chromatography and their applications to assays of drugs in serum
by J. Haginaka, J. Wakai and H. Yasuda (Nishinomiya, Japan) 163

Effect of composition of polyacrylamide gels on exclusion limits
by K. Suzuki and K. Nakazato (Kanagawaken, Japan) 173

Retention prediction based on the electrostatic model of reversed-phase ion-pair high-performance liquid chromatography: effect of pairing ion concentration
by Á. Bartha (Veszprem, Hungary) and J. Ståhlberg (Södertälje, Sweden) 181

Nature of the eddy dispersion in packed beds
by A. L. Berdichevsky and U. D. Neue (Milford, MA, U.S.A.) 189

Computer assisted internal standard selection for reversed-phase liquid chromatography
by N. E. Skelly, S. W. Barr and A. P. Zelinko (Midland, MI, U.S.A.) 199

Measurement of limiting ionic equivalent conductance by single-column ion chromatography
by Z. L. Shan, H. Xiao and P. L. Zhu (Lanzhou, China) 207

Elution orders in the separation of enantiomers by high-performance liquid chromatography with some chiral stationary phases
by N. Ôi, H. Kitahara and R. Kira (Osaka, Japan) 213

Resolution of enantiomeric lorazepam and its acyl and O-methyl derivatives and racemization kinetics of lorazepam enantiomers
by X.-L. Lu and S. K. Yang (Bethesda, MD, U.S.A.) 229

Resolution of enantiomeric triols, triol-hydroxyethylthioethers, and methoxy-triols derived from three benzo[*a*]pyrene diol-expoxides by chiral stationary phase high-performance liquid chromatography
by H. B. Weems and S. K. Yang (Bethesda, MD, U.S.A.) 239

Assay of hydroxyfarrerol in biological fluids
by A. Marzo, E. Arrigoni Martelli and G. Bruno (Rome, Italy) and E. M. Martinelli and G. Pifferi (Milan, Italy) 255

Determination of saterinone enantiomers in plasma samples with an internal standard using a Chiralcel OD column, fractionation and reversed-phase chromatography
by M. Rudolph, D. Volk and G. Schmiedel (Hamburg, F.R.G.) 263

Simultaneous separation and determination (in serum) of phenytoin and carbamazepine and their deuterated analogues by high-performance liquid chromatography–ultraviolet detection for tracer studies
by G. K. Szabo and R. J. Pylilo (Boston, MA, U.S.A.), R. J. Perchalski (Gainesville, FL, U.S.A.) and T. R. Browne (Boston, MA, U.S.A.) 271

Simultaneous determination of *p*-hydroxylated and dihydrodiol metabolites of phenytoin in urine by high-performance liquid chromatography
by G. K. Szabo, R. J. Pylilo, H. Davoudi and T. R. Browne (Boston, MA, U.S.A.) . . . 279

Automated liquid chromatographic determination of ranitidine in microliter samples of rat plasma
by A. B. Segelman (Piscataway, NJ, U.S.A.) and V. E. Adusumalli and F. H. Segelman (Cranbury, NJ, U.S.A.) 287

Analysis of carbamazepine in serum by liquid chromatography with direct sample injection and surfactant-containing eluents
by D. Bentrop, F. V. Warren, Jr., S. Schmitz and B. A. Bidlingmeyer (Milford, MA, U.S.A.) 293

Micro-scale liquid chromatographic method for the determination of bamifylline and its major metabolite in human plasma
by F. Belliardo (Turin, Italy) and C. Lucarelli (Rome, Italy) 305

Determination of phenol, *m*-, *o*- and *p*-cresol, *p*-aminophenol and *p*-nitrophenol in urine by high-performance liquid chromatography
by A. Brega (Concesio, Italy), P. Prandini and C. Amaglio (Brescia, Italy) and E. Pafumi (Concesio, Italy) 311

Rapid and sensitive detection of citrinin production during fungal fermentation using high-performance liquid chromatography
by R. B. Vail and M. J. Homann (Union, NJ, U.S.A.) 317

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)